

# Fixed points on abstract structures without the equality test

M. V. Korovina

In this paper, we present a study of the definability properties of fixed points of effective operators on abstract structures without the equality test. In particular, we prove that the Gandy theorem holds for the reals without the equality test. This provides a useful tool for dealing with recursive definitions using  $\Sigma$ -formulas.

## 1. Introduction

The aim of the paper is to present a study of the definability properties of fixed points of effective operators on abstract structures without the equality test. The question of definability of fixed points of  $\Sigma$ -operators on abstract structures with equality was first studied in [1, 6, 5]. One of the most fundamental theorems in the area is the Gandy theorem which states that the least fixed point of any positive  $\Sigma$ -operator is  $\Sigma$ -definable. This theorem allows us to treat the inductive definitions using  $\Sigma$ -formulas. The role of inductive definability as the basic principle of general computability is discussed in [9, 13]. In some cases it is natural to consider a structure in the language without equality. For example, in all effective approaches to exact real number computation via concrete representations [7, 8, 14], the equality test is undecidable. This is not surprising, because infinite amount of information should be checked in order to decide that two given numbers are equal.

Until now there has been no Gandy-type theorem known for such structures. Let us note that in all proofs of the Gandy theorem that have been known so far it is the case that, even when the definition of a  $\Sigma$ -operator does not involve equality, the resulting  $\Sigma$ -formula usually does. In this paper we show that it is possible to overcome this problem. In particular, we show that the Gandy theorem holds for the real numbers without the equality test.

The concept of  $\Sigma$ -definability is closely related to the generalised computability on an abstract structure [1, 6, 12, 15], in particular, on the real numbers [10, 11, 15].

The notions of  $\Sigma$ -definable sets or relations generalise those of computable enumerable sets of natural numbers and play a leading role in the

specification theory that is used in the higher order computation theory on abstract structures.

In this paper, we investigate definability of the least fixed points of  $\Sigma$ -operators on abstract structures without the equality test. The paper is organised as follows. In Section 2, we introduce the basic notations and definitions. We provide the background information necessary to understand the main results. Section 3 presents the Gandy theorem for structures without the equality test. In Section 4, we give an application of our result to the real numbers without the equality test. We end with a discussion of our future work.

## 2. Background

Here we introduce the basic notations and definitions. Let us consider an abstract structure  $A$  in a finite language  $\sigma_0$  without the equality test.

In order to do any kind of computation or to develop a computability theory, one has to work within a structure rich enough in information to be coded and stored. For this purpose, we extend the structure  $A$  by the set of hereditarily finite sets  $\text{HF}(A)$ .

The idea that the hereditarily finite sets over  $A$  form a natural domain of computation is quite classical and is developed in detail in [1, 6].

Note that this or very similar extensions of structures with equality are used in the theory of abstract state machines [2, 3] and in query languages for hierarchic databases [4].

We will construct the set of hereditarily finite sets over the model without equality. This structure permits us to define the natural numbers and to code and store information via formulas.

We construct the set of hereditarily finite sets,  $\text{HF}(A)$ , as follows:

1.  $\text{HF}_0(A) \doteq A$ ,  $\text{HF}_{n+1}(A) \doteq \mathcal{P}_\omega(\text{HF}_n(A)) \cup \text{HF}_n(A)$ , where  $n \in \omega$  and for every set  $B$ ,  $\mathcal{P}_\omega(B)$  is the set of all finite subsets of  $B$ .
2.  $\text{HF}(A) = \bigcup_{n \in \omega} \text{HF}_n(A)$ .

We define  $\mathbf{HF}(A)$  as the following model:

$$\mathbf{HF}(A) \doteq \langle \text{HF}(A), U, S, \sigma_0, \emptyset, \in \rangle \doteq \langle \text{HF}(A), \sigma \rangle,$$

where the constant  $\emptyset$  stands for the empty set and the binary predicate symbol  $\in$  has the set-theoretic interpretation. We also add the predicate symbols  $U$  for urelements (elements from  $A$ ) and  $S$  for sets. Let us denote  $S(\text{HF}(A)) \doteq \text{HF}(A) \setminus A$ .

The natural numbers  $0, 1, \dots$  are identified with the (finite) ordinals in  $\mathbf{HF}(A)$ , i.e.  $\emptyset, \{\emptyset, \{\emptyset\}\}, \dots$ , so in particular,  $n + 1 = n \cup \{n\}$  and the set  $\omega$  is a subset of  $\mathbf{HF}(A)$ .

We use variables subject to the following conventions:

- $r, r_1, \dots$  range over  $A$  (urelements),
- $x, y, z, s, w, f, g, \dots$  range over  $S(\mathbf{HF}(A))$  (sets),
- $n, m, l, \dots$  range over  $\omega$  (natural numbers) and
- $a, b, c, \dots$  range over  $\mathbf{HF}(A)$ .

We use the same letters to denote elements from the corresponding structures and  $\bar{r}$  to denote  $r_1, \dots, r_m$ .

The notions of a term and an atomic formula are given in the standard manner.

The set of  $\Delta_0$ -formulas is the closure of the set of atomic formulas under  $\wedge, \vee, \neg$ , and bounded quantifiers  $(\exists a \in s)$  and  $(\forall a \in s)$ , where  $(\exists a \in s) \Psi$  denotes  $\exists a(a \in s \wedge \Psi)$  and  $(\forall a \in s) \Psi$  denotes  $\forall a(a \in s \rightarrow \Psi)$ .

The set of  $\Sigma$ -formulas is the closure of the set of  $\Delta_0$  formulas under  $\wedge, \vee, (\exists a \in s), (\forall a \in s)$ , and  $\exists$ .

We are interested in  $\Sigma$ -definability of sets on  $A^n$  which can be considered as generalisation of recursive enumerability. The analogy between  $\Sigma$ -definable and recursive enumerable sets is based on the following fact. Consider the structure  $\mathbf{HF} = \langle \mathbf{HF}(\emptyset), \in \rangle$  with the hereditarily finite sets over  $\emptyset$  as its universe and membership as its only relation. In  $\mathbf{HF}$  the  $\Sigma$ -definable sets are exactly the recursively enumerable sets.

The notion of  $\Sigma$ -definability has a natural meaning also in the structure  $\mathbf{HF}(A)$ .

**Definition 1.**

1. A set  $B \subseteq \mathbf{HF}(A)$  is  $\Sigma$ -definable, if there exists a  $\Sigma$ -formula  $\Phi(a)$  such that  $b \in B \leftrightarrow \mathbf{HF}(A) \models \Phi(b)$ .
2. A function  $f : \mathbf{HF}(A) \rightarrow \mathbf{HF}(A)$  is  $\Sigma$ -definable, if there exists a  $\Sigma$ -formula  $\Phi(c, d)$  such that  $f(a) = b \leftrightarrow \mathbf{HF}(A) \models \Phi(a, b)$ .

**Lemma 1.**

1. The predicates  $R(a) \equiv a \in A$ ,  $S(a) \equiv a$  is a set, and  $n \in \omega$  are  $\Delta_0$ -definable.
2. The following predicates are  $\Delta_0$ -definable:  $x = y$ ,  $x = y \cap z$ ,  $x = y \cup z$ ,  $x = \langle y, z \rangle$ ,  $x = y \setminus z$  (recall that all variables  $x, y, z$  range over sets).
3. A function  $f : \omega^n \rightarrow \omega^m$  is computable if and only if it is  $\Sigma$ -definable.
4. Let  $\text{Fun}(g)$  mean that  $g$  is a finite function

$$g = \{ \langle x, y \rangle \mid \text{for every } x \text{ there exists a unique } y \}.$$

Then the predicate  $\text{Fun}(g)$  is  $\Delta_0$ -definable.

5. If  $\mathbf{HF}(A) \models \text{Fun}(g)$  then the domain of  $g$ , denoted by  $\delta_g$ , is  $\Delta_0$ -definable.

**Proof.** Proofs of all properties are straightforward except (3) which can be found in [6].  $\square$

For finite functions  $\text{Fun}(f)$ , let us denote  $f(x) = y$  if  $\langle x, y \rangle \in f$ .

The following proposition states that we have a full collection on  $\mathbf{HF}(A)$ .

**Proposition 1.** (Collection.) *For every formula  $\Phi$  the following holds: if  $\mathbf{HF}(A) \models (\forall a \in x) \exists b \Phi(a, b)$ , then there is a set  $z$  such that*

$$\mathbf{HF}(A) \models (\forall a \in x) (\exists b \in z) \Phi(a, b) \wedge (\forall b \in z) (\exists a \in x) \Phi(a, b).$$

**Proof.** This follows from the definition of  $\mathbf{HF}(A)$ . Indeed, if  $x \in \mathbf{HF}(A)$  consists of  $k$  elements  $a_1, \dots, a_k$  and for each of these  $a_i$  there is an  $b_i$  such that  $\Phi(a_i, b_i)$  holds. Then all  $b_1, \dots, b_k$  occur in  $\mathbf{HF}_n(A)$  for some element  $n$ , hence  $\{b_1, \dots, b_k\} \in \mathbf{HF}_{n+1}(A)$ .  $\square$

### 3. The least fixed points of effective operators

Now we recall the notion of a  $\Sigma$ -operator and prove the Gandy theorem for structures without the equality test.

Let  $\Phi(a_1, \dots, a_n, P)$  be a  $\Sigma$ -formula, where  $P$  occurs positively in  $\Phi$  and the arity of  $P$  is equal to  $n$ .

We think of  $\Phi$  as defining a  $\Sigma$ -operator  $\Gamma : \mathcal{P}(\mathbf{HF}(A)^n) \rightarrow \mathcal{P}(\mathbf{HF}(A)^n)$  given by

$$\Gamma(Q) = \{ \bar{a} \mid (\mathbf{HF}(A), Q) \models \Phi(\bar{a}, P) \},$$

where for every set  $B$ ,  $\mathcal{P}(B)$  is the set of all subsets of  $B$ .

Since the predicate symbol  $P$  occurs only positively, we have that the corresponding operator  $\Gamma$  is monotone, i.e. for any sets  $A \subseteq B$  implies  $\Gamma(A) \subseteq \Gamma(B)$ .

By monotonicity, the operator  $\Gamma$  has the least (w.r.t. inclusion) fixed point which can be described as follows.

We start from the empty set and apply the operator  $\Gamma$  until we reach the fixed point:

$$\Gamma^0 = \emptyset, \quad \Gamma^{\beta+1} = \Gamma(\Gamma^\beta), \quad \Gamma^\gamma = \bigcup_{\beta < \gamma} \Gamma^\beta, \quad (1)$$

where  $\gamma$  is a limit ordinal.

One can easily check that the sets  $\Gamma^\beta$  form an increasing chain of sets:  $\Gamma^0 \subseteq \Gamma^1 \subseteq \dots$ . By set-theoretical reasons, there exists the least ordinal  $\gamma$  such that  $\Gamma(\Gamma^\gamma) = \Gamma^\gamma$ . This  $\Gamma^\gamma$  is the least fixed point of the given operator  $\Gamma$ .

In order to study the least fixed points of arbitrary  $\Sigma$ -operators (without the equality test), we first consider  $\Sigma$ -operators of the type

$$\Gamma : \mathcal{P}(S(\mathbf{HF}(A))^n) \rightarrow \mathcal{P}(S(\mathbf{HF}(A))^n).$$

Then we will show how the least fixed points of arbitrary  $\Sigma$ -operators can be constructed using the least fixed points of such operators. Note that, as  $S(\mathbf{HF}(A))$  is closed under pairing,  $S(\mathbf{HF}(A))^n \subseteq S(\mathbf{HF}(A))$  for  $n > 0$ . Moreover,  $S(\mathbf{HF}(A))^n$  is a  $\Sigma$ -definable subset of  $\mathbf{HF}(A)$ . So, without loss of generality, we can consider the case  $n = 1$ .

Let us formulate some properties of  $\Sigma$ -operators which we will use below. The following proposition states that each element from the value of a  $\Sigma$ -operator on a  $\Sigma$ -set can be obtained as an element of the value of this operator on a finite subset of the set.

**Proposition 2.** *If  $Q$  is a  $\Sigma$ -definable subset of  $S(\mathbf{HF}(A))$  and  $w \in \Gamma(Q)$ , then there exists  $p \in S(\mathbf{HF}(A))$  such that  $p \subseteq Q$  and  $w \in \Gamma(p)$ .*

**Proof.** We prove the proposition for the more general case where we allow parameters from  $S(\mathbf{HF}(A))$  to occur into the formula defining our operator.

Let  $\Phi(\bar{b}, x, P)$  be a  $\Sigma$ -formula defining our operator  $\Gamma$ , where  $\bar{b} = b_1, \dots, b_n$  are parameters from  $S(\mathbf{HF}(A))$ . And let  $Q$  be a  $\Sigma$ -definable subset of  $S(\mathbf{HF}(A))$  and  $w \in \Gamma(Q)$ . We need to prove that there exists  $p \in S(\mathbf{HF}(A))$  such that  $p \subseteq Q$  and  $w \in \Gamma(p)$ .

We prove the claim by induction on the structure of  $\Phi$ .

If  $\Phi(\bar{b}, x, P) \equiv P(x)$  and  $(\mathbf{HF}(A), Q) \models P(w)$ , then the set  $p \equiv \{w\}$  is a required one.

If  $\Phi$  is an atomic formula which does not contain  $P$ , then the set  $p \equiv \emptyset$  is a required one.

For the induction step, let us consider all possible cases.

1. Suppose  $\Phi(\bar{b}, x, P) \equiv (\forall a \in b_j) \Psi(a, \bar{b}, x, P)$  and

$$(\mathbf{HF}(A), Q) \models (\forall a \in b_j) \Psi(a, \bar{b}, w, P).$$

By induction hypothesis,

$$(\mathbf{HF}(A), Q) \models (\forall a \in b_j) \exists s (\Psi(a, \bar{b}, w, P))_{t \in s}^{P(t)} \wedge s \subseteq Q.$$

Using Proposition 1, we find an element  $q$  such that

$$\begin{aligned} (\mathbf{HF}(A), Q) \models & (\forall a \in b_j) (\exists s \in q) \left( (\Psi(a, \bar{b}, w, P))_{t \in s}^{P(t)} \wedge s \subseteq Q \right) \wedge \\ & (\forall s \in q) (\exists a \in b_j) \left( (\Psi(a, \bar{b}, w, P))_{t \in s}^{P(t)} \wedge s \subseteq Q \right). \end{aligned}$$

Let  $p \equiv \cup q$ .

By definition, for all  $a \in b_j$  there exists  $s \subseteq p$  such that

$$(\mathbf{HF}(A), s) \models (\Psi(a, \bar{b}, w, P))_{t \in s}^{P(t)}.$$

So we have

$$(\mathbf{HF}(A), p) \models \Psi(a, \bar{b}, w, P) \text{ for all } a \in b_j.$$

In other words,

$$(\mathbf{HF}(A), p) \models (\forall a \in b_j) \Psi(a, \bar{b}, x, P).$$

By construction, the set  $p$  is a required one.

2. The case  $\Phi(\bar{b}, x, P) \equiv (\exists a \in b_j) \Psi(a, \bar{b}, x, P)$  is similar to the case above.

3. Suppose  $\Phi(\bar{b}, x, P) \equiv \exists a \Psi(a, \bar{b}, x, P)$  and

$$(\mathbf{HF}(A), Q) \models \Psi(b', \bar{b}, w, P).$$

By induction hypothesis, there exists  $p_0 \subseteq Q$  such that  $p_0 \in S(\mathbf{HF}(A))$  and

$$(\mathbf{HF}(A), p_0) \models \Psi(b', \bar{b}, w, P).$$

The set  $p \equiv p_0$  is a required one.

4. Suppose  $\Phi(\bar{b}, x, P) \equiv \Psi_1(\bar{b}, x, P) \wedge \Psi_2(\bar{b}, x, P)$  and

$$(\mathbf{HF}(A), Q) \models \Psi_1(\bar{b}, w, P) \wedge \Psi_2(\bar{b}, w, P).$$

By induction hypothesis, there exist  $p_1 \subseteq Q$  and  $p_2 \subseteq Q$  such that  $p_1 \in S(\mathbf{HF}(A))$ ,  $p_2 \in S(\mathbf{HF}(A))$  and

$$(\mathbf{HF}(A), p_1) \models \Psi_1(\bar{b}, w, P)$$

and

$$(\mathbf{HF}(A), p_2) \models \Psi_2(\bar{b}, w, P).$$

The set  $p \equiv p_1 \cup p_2$  is a required one.

5. The case  $\Phi(\bar{b}, x, P) \equiv \Psi_1(\bar{b}, x, P) \vee \Psi_2(\bar{b}, x, P)$  is similar to the case above.  $\square$

**Proposition 3.** *Let  $\Gamma : \mathcal{P}(S(\mathbf{HF}(A))) \rightarrow \mathcal{P}(S(\mathbf{HF}(A)))$  be a  $\Sigma$ -operator. The relation  $x \in \Gamma(y)$  is  $\Sigma$ -definable.*

**Proof.** Let  $\Phi(z, P)$  be a  $\Sigma$ -formula which defines the operator  $\Gamma$ . Suppose  $x \in \Gamma(y)$ . By definition,

$$x \in \{z \mid (\mathbf{HF}(A), y) \models \Phi(z, P)\}.$$

It means that

$$(\mathbf{HF}(A), y) \models \Phi(x, P).$$

So we have

$$(\mathbf{HF}(A)) \models (\Phi(x, P))_{t \in y}^{P(t)}.$$

It is easy to see that the relation  $x \in \Gamma(y)$  is defined by  $\Sigma$ -formula  $\Phi(x, P)_{t \in y}^{P(t)}$ .  
□

Now we are ready to prove the Gandy theorem for  $\Sigma$ -operators of the type

$$\Gamma : \mathcal{P}(S(\mathbf{HF}(A))) \rightarrow \mathcal{P}(S(\mathbf{HF}(A))).$$

**Theorem 1.** *Let  $\Gamma : \mathcal{P}(S(\mathbf{HF}(A))) \rightarrow \mathcal{P}(S(\mathbf{HF}(A)))$  be a  $\Sigma$ -definable operator. Then the least fixed-point of  $\Gamma$  is  $\Sigma$ -definable.*

**Proof.** We will prove that the least fixed point of the operator  $\Gamma$  is  $\Gamma^\omega$ , where  $\Gamma^\omega$  is defined as follows:  $\Gamma^0 = \emptyset$ ,  $\Gamma^n = \Gamma(\Gamma^{n-1})$  for a finite ordinal  $n$ , and  $\Gamma^\omega = \bigcup_{m < \omega} \Gamma^m$ .

Let us show  $\Sigma$ -definability of  $\Gamma^n$  for every finite ordinal  $n$ .

For this purpose, we introduce the following family of finite functions:

$$\begin{aligned} X_0 &= \{ \langle \emptyset, \emptyset \rangle \}, \\ X_n &= \{ f \mid Fun(f) \text{ and } \delta_f = n + 1, f(0) = \emptyset, f \text{ is monotonic} \\ &\quad \text{and for any } m \leq n \text{ the following is true: } f(m) \subseteq \bigcup_{l < m} \Gamma(f(l)) \} \end{aligned}$$

where  $n > 0$ , and  $S_j$  is the domain of the function.

From the definitions  $X_n$  and  $\Gamma$ , it follows that  $X_n$  is  $\Sigma$ -definable for all  $n \in \omega$ , moreover there exists a  $\Sigma$ -formula  $\psi(n, x)$  such that

$$\mathbf{HF}(A) \models \psi(n, x) \leftrightarrow x \in X_n.$$

Below we will use the following useful properties of the families  $X_n$ :

1. Let  $w$  be a finite subset of  $X_n$ . Let us define  $f^*(m) = \bigcup_{f \in w} f(m)$  for all  $m \leq n$ . Then  $f^* \in X_n$ .

2. If  $f \in X_n$  and  $m \leq n$ . Then  $f \upharpoonright (m+1) \in X_m$ .
3. Let  $f \in X_m$  and  $m \leq n$ .

Define a function

$$f^*(l) = \begin{cases} f(l), & \text{if } l \leq m \\ f(m), & \text{if } m < l \leq n. \end{cases}$$

Then  $f^* \in X_n$ .

4. Let  $f \in X_n$  and  $b \in \Gamma(f(m))$ , where  $m \leq n$ .

Define a function

$$f^*(l) = \begin{cases} f(l), & \text{if } l \leq n \\ \{b\}, & \text{if } l = n+1. \end{cases}$$

Then  $f^* \in X_{n+1}$ .

Using these properties, let us show that:

$$x \in \Gamma^n \text{ iff } \mathbf{HF}(A) \models \exists f (f \in X_n \wedge x \in f(n)) \quad (2)$$

by induction on  $n$ . For  $n = 0$  we have  $\Gamma^0 = \emptyset$  and therefore (2) holds.

Assume that (2) holds for  $n$ . Let us prove that (2) holds for  $n+1$ .

To prove from left to right, let us consider  $x \in \Gamma^{n+1} = \Gamma(\Gamma^n)$ . By induction hypothesis, we have that  $x_1 \in \Gamma^n$  iff  $\exists g (g \in X_n \wedge x_1 \in g(n))$ . So the set  $\Gamma^n$  is  $\Sigma$ -definable. By Proposition 2, it follows that there exists  $y \in S(\mathbf{HF}(A))$  such that  $y \subseteq \Gamma^n$  and  $x \in \Gamma(y)$ .

By induction hypothesis and the condition  $y \subseteq \Gamma^n$ ,

$$\mathbf{HF}(A) \models (\forall z \in y) \exists g (g \in X_n \wedge z \in g(n)).$$

Using Proposition 1, we find an element  $w$  such that

$$\begin{aligned} \mathbf{HF}(A) \models & (\forall z \in y) (\exists g \in w) (g \in X_n \wedge z \in g(n)) \wedge \\ & (\forall g \in w) (\exists z \in y) (g \in X_n \wedge z \in g(n)). \end{aligned}$$

Starting from the finite subset  $w \subseteq X_n$ , we define the function  $g_0$  as follows:

$$g_0(l) = \cup_{g \in w} g(l), \quad l \leq n.$$

By Property (1) of  $X_n$  which is mentioned above,  $g_0 \in X_n$ . It is easy to check the following inclusion:  $y \subseteq g_0(n)$ . Indeed, if  $z \in y$ , then there exists  $g \in w$  such that  $z \in g(n) \subseteq g_0(n)$ .

Define a function



$$f(l) = \begin{cases} g_0(l), & \text{if } l \leq n \\ \{x\}, & \text{if } l = n + 1. \end{cases}$$

From Property (4) of  $X_n$ , it follows that  $f \in X_{n+1}$  and, moreover,  $x \in f(n+1)$  holds by the definition of  $f$ . So  $f$  is a required one.

To prove from right to left, let us suppose that there exists  $f$  such that  $(f \in X_{n+1} \wedge x \in f(n+1))$ . By the definition of  $X_{n+1}$ ,  $x \in \Gamma(f(m))$  for some  $m \leq n$ .

Let us check the inclusion :  $f(m) \subseteq \Gamma^m$ . For this purpose we consider  $f_1 = f \upharpoonright (m+1)$ . From Property (2) of  $X_m$ , it follows that  $f_1 \in X_m$ . So, for all  $y \in f_1(m)$  we have  $\mathbf{HF}(A) \models \exists f (f \in X_m \wedge y \in f(m))$ . By induction, it means that  $f_1(m) = f(m) \subseteq \Gamma^m$ .

The operator  $\Gamma$  is monotone, so we have

$$x \in \Gamma(f(m)) \subseteq \Gamma(\Gamma^m) \subseteq \bigcup_{m < n+1} \Gamma(\Gamma^m) = \Gamma^{n+1}.$$

Thus we have proven that  $\Gamma^n$  is  $\Sigma$ -definable for all  $n \in \omega$ . Consequently,

$$x \in \Gamma^\omega \leftrightarrow \exists n \exists f (f \in X_n \wedge x \in f(n)) \quad (3)$$

is  $\Sigma$ -definable.

To check that  $\Gamma^\omega$  is a fixed point, i.e.  $\Gamma(\Gamma^\omega) \subseteq \Gamma^\omega$  let us consider  $x \in \Gamma(\Gamma^\omega)$ . From (3) it follows that  $\Gamma^\omega$  is  $\Sigma$ -definable. From Proposition 2 it follows that there exists  $y \in S(\mathbf{HF}(A))$  such that  $y \subseteq \Gamma^\omega$  and  $x \in \Gamma(y)$ . It is easy to check that  $y \subseteq \Gamma^m$  for some  $m \in \omega$ . From this we have that  $x \in \Gamma(\Gamma^m) \subseteq \Gamma^\omega$ . By monotonicity of  $\Gamma$ , the set  $\Gamma^\omega$  is the least fixed point. So the least fixed point of the operator  $\Gamma$  is  $\Sigma$ -definable.  $\square$

Now we consider arbitrary  $\Sigma$ -operators on the structure  $A$  without the equality test.

**Theorem 2.** *Let  $\Gamma : \mathcal{P}(\mathbf{HF}(A)^n) \rightarrow \mathcal{P}(\mathbf{HF}(A)^n)$  be an arbitrary  $\Sigma$ -operator. Then the least fixed-point of  $\Gamma$  is  $\Sigma$ -definable.*

**Proof.**

Without loss of generality, let us consider the case  $n = 1$ . For simplicity of notation, we will give the construction only for that case, since the main ideas are already contained here. Let  $\Phi(r, P)$  define the operator  $\Gamma$ . We construct a new  $\Sigma$ -operator  $F : \mathcal{P}(S(\mathbf{HF}(A))) \rightarrow \mathcal{P}(S(\mathbf{HF}(A)))$  such that

$$r \in \Gamma^n \longleftrightarrow \exists x (x \in F^n \wedge r \in x).$$

For this purpose we define the following formula with a new unary predicate symbol  $Q$ :

$$\Psi(x, Q) = (\forall r \in x) (\Phi(r, P))_{\exists y Q(y) \wedge t \in y}^{P(t)}.$$

It is easy to see that  $\Psi$  induces a  $\Sigma$ -operator  $F$  given by

$$F(D) = \{x \mid (\mathbf{HF}(A), D) \models \Psi(x, Q)\}.$$

Let us show that

$$r \in \Gamma^n \leftrightarrow \exists x (x \in F^n \wedge r \in x) \quad (4)$$

by induction on  $n$ . For  $n = 0$  we have  $\Gamma^n = F^n = \emptyset$  and therefore (4) holds.

Assume that (4) holds for  $n$ . Let us prove that (4) holds for  $n + 1$ . In other words, we need to prove that

$$\begin{aligned} (\mathbf{HF}(A), \Gamma^n) \models \Phi(r, P) &\leftrightarrow \\ (\mathbf{HF}(A), F^n) \models \exists x \left( r \in x \wedge (\forall r' \in x) (\Phi(r', P))_{\exists y Q(y) \wedge t \in y}^{P(t)} \right). \end{aligned}$$

Since the first formula does not contain  $Q$  and the second formula does not contain  $P$ , it is sufficient to consider one structure  $(\mathbf{HF}(A), \Gamma^n, F^n)$  and prove that

$$\begin{aligned} (\mathbf{HF}(A), \Gamma^n, F^n) \models \Phi(r, P) &\leftrightarrow \\ (\mathbf{HF}(A), \Gamma^n, F^n) \models \exists x \left( r \in x \wedge (\forall r' \in x) (\Phi(r', P))_{\exists y Q(y) \wedge t \in y}^{P(t)} \right). \end{aligned}$$

To prove from left to right, let us consider  $r \in \mathbf{HF}(A)$  such that

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models \Phi(r, P).$$

Consider the formula  $(\Phi(r, P))_{\exists y Q(y) \wedge t \in y}^{P(t)}$ . Then by induction hypothesis we have that

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models \forall r' (P(r') \leftrightarrow \exists x (x \in Q \wedge r' \in x)) \quad (5)$$

and therefore (by replacement lemma) we have

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models (\Phi(r, P))_{\exists y Q(y) \wedge t \in y}^{P(t)}.$$

Now it is easy to check that

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models \exists x \left( r \in x \wedge (\forall r' \in x) (\Phi(r', P))_{\exists y Q(y) \wedge t \in y}^{P(t)} \right)$$

taking  $x = \{r\}$ .

To prove from right to left, let us consider  $r \in \mathbf{HF}(A)$  such that

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models \exists x \left( r \in x \wedge (\forall r' \in x) (\Phi(r', P))_{\exists y Q(y) \wedge t \in y}^{P(t)} \right).$$

From this we have that

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models (\Phi(r, P))_{\exists y Q(y) \wedge t \in y}^{P(t)}$$

and from (5) (by the replacement lemma) we obtain that

$$(\mathbf{HF}(A), \Gamma^n, F^n) \models \Phi(r, P).$$

Now, from Theorem 1, it follows that the least fixed point of the operator  $F$  is  $\Sigma$ -definable and therefore the least fixed point of the operator  $\Gamma$  is also  $\Sigma$ -definable. □

#### 4. The least fixed points of effective operators on the real numbers without the equality test

In this section, we consider the standard model of the real numbers  $\langle \mathbb{R}, 0, 1, +, \cdot, -, < \rangle$ , denoted also by  $\mathbb{R}$ , where  $+$ ,  $\cdot$  and  $-$  are regarded as the usual arithmetic operations on the reals. We use the language of strictly ordered rings, so the predicate  $<$  occurs positively in formulas. This allows us to consider  $\Sigma$ -definability as generalisation of computable enumerability. Indeed, in all effective approaches to exact real number computation via concrete representations, we need only finite amount of information in order to show that one number is less than another. The following is an immediate corollary of Theorem 2.

**Corollary 1.** *Let  $\Gamma : \mathcal{P}(\mathbf{HF}(\mathbb{R})^n) \rightarrow \mathcal{P}(\mathbf{HF}(\mathbb{R})^n)$  be an arbitrary  $\Sigma$ -operator. Then the least fixed-point of  $\Gamma$  is  $\Sigma$ -definable.*

#### 5. Future work

One of the applications of the Gandy theorem in the case of structures with equality is that it allows us to define universal  $\Sigma$ -predicates. It leads to a topological characterisation of  $\Sigma$ -relations on  $\mathbb{R}$ . Thus the sets  $B \subseteq \mathbb{R}^n$  that are  $\Sigma$ -definable in  $\mathbf{HF}(\mathbb{R})$  with equality are exactly the effective unions of semialgebraic sets.

We think that the Gandy theorem can be used in this way for the structures without equality, but for this we need more evolved arguments. Also we think that it is possible to show that the sets  $B \subseteq \mathbb{R}^n$  that are  $\Sigma$ -definable in  $\mathbf{HF}(\mathbb{R})$  without equality are exactly the effective unions of open semialgebraic sets.

## References

- [1] Barwise J. *Admissible Sets and Structures*. — Berlin, Springer-Verlag, 1975.
- [2] Blass A., Gurevich Y., Shelah S. Choiceless polynomial time // *Pure and Applied logic*. — 1999. — N 100. — P. 141–187.
- [3] Blass A., Gurevich Y. Background, reserve and Gandy machines // *Proc. CSL'2000*. — *Lect. Notes Comput. Sci.* — 2000. — Vol. 1862. — P. 1–17.
- [4] Dahlhaus E., Makowsky J. A. Query languages for hierarchic databases // *Information and Computation*. — 1992. — N 101. — P. 1–32.
- [5] Gandy R. Inductive definitions // *Generalized Recursion Theory*. — Amsterdam: North-Holland, 1974. — P. 265–300.
- [6] Ershov Yu. L. *Definability and Computability*. — N.-Y.: Plenum Press, 1996.
- [7] Freedman H., Ko K. Computational complexity of real functions // *Theor. Comput. Sci.* — 1982. — Vol. 20. — P. 323–352.
- [8] Grzegorzczak A. On the definitions of computable real continuous functions // *Fund. Math.* — 1957. — N 44. — P. 61–71.
- [9] Hinman P. G. Recursion on abstract structures // *Handbook of Computability Theory*. — Elsevier, 1999. — P. 317–359.
- [10] Korovina M., Kudinov O. Characteristic Properties of Majorant-Computability over the Reals // *Proc. of CSL'98*. — *Lect. Notes Comput. Sci.* — 1999. — Vol. 1584. — P. 188–204.
- [11] Korovina M., Kudinov O. Semantic characterisations of second-order computability over the real numbers // *Proc. of CSL'01*. — *Lect. Notes Comput. Sci.* — 2001. — Vol. 2142. — P. 160–173.
- [12] Moschovakis Y. N. Abstract first order computability // *Trans. Amer. Math. Soc.* — 1969. — Vol. 138. — P. 427–464.
- [13] Moschovakis Y. N. *Elementary Induction on Abstract Structure*. — North-Holland, 1974.
- [14] Pour-El M. B., Richards J. I., *Computability in Analysis and Physics*. — Springer-Verlag, 1988.
- [15] Tucker J. V., Zucker J. I., *Computable functions and semicomputable sets on many-sorted algebras* // *Handbook of Logic in Computer Science*. — Oxford: Oxford University Press, 2000. — Vol. 5. — P. 397–525.